

1 **Generation of bioactive peptides during food processing**
2

3
4 Fidel Toldrá^{1,3}, Milagro Reig², M-Concepción Aristoy¹ and Leticia Mora¹
5

6
7 ¹*Instituto de Agroquímica y Tecnología de Alimentos (CSIC), Paterna (Valencia), Spain*
8 and ²*Instituto de Ingeniería de Alimentos para el Desarrollo, Universidad Politécnica*
9 *de Valencia, Spain*
10

11
12
13
14
15 ³Corresponding author: Fidel Toldrá
16

17 e-mail: ftoldra@iata.csic.es

18 Tel: +34 963636301

19 Fax: +34 963900022
20

21
22
23
24
25
26 Running title: **Generation of bioactive peptides during food processing**
27
28

Abstract

Large amounts of peptides are naturally generated in foods through the proteolysis phenomena taking place during processing. Such proteolysis is carried out either by endogenous enzymes in ripened foods or by the combined action of endogenous and microbial enzymes when fermented. Food proteins can also be isolated and hydrolysed by peptidases to produce hydrolysates. Endo-peptidases act first followed by the successive action of exo-peptidases (mainly, tri- and di-peptidylpeptidases, aminopeptidases and carboxypeptidases). The generated peptides may be further hydrolysed through the gastrointestinal digestion resulting in a pool of peptides with different sequences and lengths, some of them with relevant bioactivity. However, these peptides should be absorbed intact through the intestinal barrier and reach the blood stream to exert their physiological action. This manuscript is reporting the enzymatic routes and strategies followed for the generation of bioactive peptides.

Keywords: proteolysis, peptides, bioactive peptides, proteomics, enzymes, peptidases, exo-peptidases

1 Introduction

Proteins are one of the main components in foods, especially in those of animal origin such as meat, fish, milk or eggs. Proteins exert nutritional, functional and biological properties which are frequently affected by the technology used in food processing. Main causes of alteration of proteins during processing are pH changes like acidification, chemical treatments such as acylation, glycosylation, and phosphorylation, heat treatments and fermentation (Pihlanto & Korhonen, 2003). These changes can be responsible for positive aspects such as the improvement in final textural/organoleptic characteristics, better stability of the product, or the generation of bioactive peptides, although some negative aspects such as the modification of one or several amino acids or the generation of allergenic compounds can affect the product. Changes occurred during specific food processes such as curing or fermentation in cheese, wine or dry-cured meats have been widely described as a source of bioactive peptides (Corrêa et al., 2014; Mohanty, Mohapatra, Misra & Sahu, 2016; Mora, Escudero, Arihara & Toldrá, 2015). Other key mechanisms to obtain bioactive peptides are the hydrolysis using controlled and commercial peptidases or microorganisms mainly used to take advantage of food by-products (Ryder, Bekhit, McConnell & Carne, 2016). Finally, the gastrointestinal (GI) digestion due to the action of salivary, stomachal, intestinal and pancreatic enzymes constitute the final hydrolysis step in generating bioactive peptides (Capriotti, Caruso, Cavaliere, Samperi et al., 2015; Pepe et al., 2016). The role of food proteins as a source of bioactive peptides has been widely described in recent studies (Li-Chan, 2015; Oseguera-Toledo, González de Mejía, Reynoso-Camacho, Cardador-Martínez & Amaya-Llano, 2014; Lassoued et al., 2015). In this respect, the bioactive peptides are inactive when they are taking part of the parent protein, but turn active when released due to the action of enzymes during food

processing or GI digestion. Once released, the bioactive peptides may provide different functions that can be reproduced *in vitro* with biochemical assays or *in vivo* in cell or animal models and humans. Different open access databases report the bioactive peptides that are being discovered including data about their main chemical and structural characteristics, IC₅₀, protein of origin, and references. Most studied biological functions to date according to the results reported by BIOPEP database (<http://www.uwm.edu.pl/biochemia/index.php/pl/biopep>) have been ACE-inhibitory activity, antioxidant activity, antimicrobial activity, opioid activity, immunomodulating and antithrombotic activities.

Numerous studies in the literature have reported bioactivities derived from food protein sources. First discoveries of food derived peptides were from milk-based products which have been extensively studied in relation to their potential health promoting effects in humans (Nongonierma and FitzGerald, 2016). Also meat protein is considered a good source to obtain bioactive peptides due to the high quality of its proteins which has been widely described (Escudero, Sentandreu, Arihara & Toldrá, 2010; Udenigwe & Howard, 2013;) together with fish proteins (Ferraro, Carvalho, Piccirillo, Santos, Castro, Pintado, 2013). The use of by-products obtained from protein sources such as slaughterhouses, fisheries, olivemill wastewater, cheese whey, winery sludge, citrus peel, etc is done under controlled enzymatic conditions which permits the control of the hydrolysis and the generated peptides (Mora, Reig & Toldrá, 2014; Ryder, Bekhit, McConnell & Carne, 2016). Egg proteins, soybean proteins, or peanut proteins have also been extensively studied protein hydrolysates (Ji, Sun, Zhao, Xiong & Sun, 2014; Tanzadehpanah, Asoodeh & Chamani, 2012; De Oliveira, Corrêa, Coletto, Daroit, Cladera-Olivera & Brandelli, 2015).

The effect of natural bioactive peptides on health by preventing infection and diseases is of great interest nowadays due to the severe toxic side-effects that have been described to be caused by the use of synthetic peptides and drugs in the treatment and prevention of numerous diseases. Also the economic impact on health care in future years due to the effect of bad habits and ageing of population could be decreased by proportionally increasing the development and use of bioactive peptides.

2. Main characteristics of peptides exerting biological activities

Data about the characteristics of bioactive peptides such as their length, amino acid composition or structural conformation results very useful in the identification and characterisation of novel active sequences, especially when empirical strategies are used in the detection of bioactive peptides. These characteristics are only well-known in the most studied activities as the lack of information about peptide sequences identified in minor bioactivities as well as the ideal conditions of proteolysis makes more difficult the standardisation of the peptides characteristics.

2.1. ACE-inhibitory activity

Angiotensin I-converting enzyme (ACE) is a key enzyme influencing the regulation of blood pressure. ACE is a central component of the renin-angiotensin system, and converts angiotensin I into the potent vasoconstrictor angiotensin II. It is also well-known to degrade the vasodilative bradykinin in the kinin–kallikrein system. For these reasons, the inhibition of ACE enzyme is of high interest in the search of antihypertensive peptides (Escudero, Mora & Toldrá, 2014).

ACE is an exopeptidase with an ability to cleave dipeptides from the C-terminal of peptides. It is a chloride-activated zinc metallopeptidase and it is assumed that the function of the anion activation in ACE provides high *in vitro* substrate specificity.

Studies with different peptide inhibitors showed that binding to ACE is strongly influenced by the C-terminal tripeptide sequence of the substrate. In fact, main inhibitors of ACE enzyme show hydrophobic amino acid residues at each of the three C-terminal positions with proline, lysine or arginine as C-terminal amino acids (Fernández, Benito, Martín, Casquete, Córdoba & Córdoba, 2016; Gu, Majumder & Wu, 2011).

2.2. Antioxidant activity

The antioxidant activity showed by peptides is classified into two groups depending on the basis of the chemical reactions involved: methods based on hydrogen atom transfer (HAT) and methods based on electron transfer (ET) (Huang, Ou, & Prior, 2005). The HAT-based assays evaluate the ability of a peptide to reduce free radicals by hydrogen donation in a competitive reaction. The *in vitro* assays used for its measurement are oxygen radical absorbance capacity assay (ORAC), total radical trapping antioxidant parameter (TRAP) and β -carotene bleaching assay. ET-based assays evaluate the ability of a potential antioxidant to transfer one electron to reduce an oxidant, so these reactions are pH dependent. The assays ABTS radical scavenging assay, ferric-reducing antioxidant power, and DPPH radical scavenging activity are used for its measurement (Huang, Ou & Prior, 2005; McDonald-Wicks, Wood, & Garg, 2006).

According to Liu, Xing, Fu, Zhou and Zhang (2016), most of the antioxidant peptides have between 4–16 amino acids, with molecular mass of about 400–2000 Da. The molecular weight affects the used routes to reach target sites and the capacity to suffer additional digestion by GI enzymes which could increase the antioxidant capacity *in vivo* (Li, Le, Shi & Shrestha, 2004). Peptides showing Pro residues have been described to be more resistant to further degradation by digestive enzymes (Fitzgerald & Meisel, 2000).

The type of amino acid plays an important role in determining the antioxidant activity of the peptides. In this respect, aromatic amino acids such as Tyr, His, Trp, and Phe can donate protons contributing to the radical-scavenging properties (Rajapakse, Mendis, Jung, Je & Kim, 2005). On the other hand, the hydrophobic amino acids have been described to be able to increase the presence of peptides at the water-lipid interface and then access to scavenge free radicals from the lipid phase (Ranathunga, Rajapakse & Kim, 2006). Finally, acidic amino acids utilise carbonyl and amino groups in the side chain which function as chelators of metal ions (Suetsuna, Ukeda & Ochi, 2000).

2.3. Antimicrobial activity

Antimicrobial peptides generated from dietary proteins show several characteristic properties. They are relatively small (20-46 amino acid residues), basic (lysine- or arginine-rich), and amphipathic. The mechanism of action of these antimicrobial peptides is still not well-known but it is believed that their effectivity depends on their capacity to form channels or pores within the microbial membrane impairing the possibility for anabolic processes (Castellano, Mora, Escudero, Vignolo, Aznar & Toldrá, 2016). Antibacterial peptides are usually described as long chains, which can adopt an α -helical linear or circular structure organized in a β -sheet which is essential against microorganisms.

2.4. Opioid activity

Exorphin is the name of those opioid peptides derived from exogenous proteins. The classic opioid peptides show the N-terminal tetra-peptide sequence Tyr-Gly-Gly-Phe, although many opioid active peptides have been described containing the N-terminal sequence Tyr-Pro. So, many opioid peptides isolated from mammalian and amphibian

sources share a common short sequence fragment with Tyr at the N-terminal (except a-casein opioids) separated from a Phe residue or the aromatic tyrosine by one or two amino acids (Stefanucci, Mollica, Macedonio, Zengin, Ahmed & Novellino, 2017). The C-terminal sequences of these peptides vary substantially both in sequence and length, but the described structural motif fits into the binding site of opioid receptors. The negative potential of the tyrosine amino acid is essential for opioid activity and the removal of Tyr residue from the active peptide results in the absence of activity (Guesdon, Pichon & Tomé, 2006).

It has been described that in *in vitro* assays, exorphins resulted from one hundred to one thousand times more potent than endogenous opioid peptides. On the other hand, some exogenous opioid peptides are active after oral administration, in which none of the exorphins were active. One of the reasons could be that Tyr-Pro sequence is more resistant to enzymatic GI digestion than the characteristic sequences for endogenous opioid peptides.

2.5. Immunomodulating activity

Immunomodulating activity has been especially identified in peptides derived from milk and milk products. These studies show that their length can be very different comprising from 2 to 64 amino acids, although those smaller than 3000 Da are the most abundant. The most repeated amino acids in the active sequences are Pro and Glu, with Tyr and Lys in the N-terminal and C-terminal extremes, respectively, and Arg at both extremes. Also, their charges differ widely at physiological pH, between 7 and 8, being mainly of hydrophilic character (Reyes-Díaz, González-Córdova, Hernández-Mendoza & Vallejo-Córdova, 2016).

3. Mechanisms for proteolysis phenomena

Proteins are hydrolysed step-wise by peptidases within the food, from initial proteins and polypeptides down to sequences with just a few amino acids. This proteolysis may take place within the food during processing by endogenous peptidases and/or by microbial peptidases in fermented foods as schematised in figure 1. Such microorganisms have a variety of enzymes which are able to hydrolyse proteins, carbohydrates and lipids (Flores & Toldrá, 2011).

The result of the combined action of endo and exo-peptidases is an accumulation of small peptides and free amino acids in foods. As it has been previously described, some of the released peptides may be bioactive if showing the adequate length and sequence of residues. An scheme on how proteolysis proceeds in foods by endogenous or microbial peptidases generating small amounts of bioactive peptides is shown in figure 2. Proteins may be also isolated from foods and hydrolysed with commercial peptidases releasing large amounts of bioactive peptides (see figure 2). Of course, the generated peptides must be ingested, subject to gastrointestinal digestion and absorbed intact through the intestinal barrier and reach the blood stream to exert their physiological action (Gallego, Grootaert, Mora, Aristoy, Van Camp & Toldrá, 2015).

The application of peptidomics tools allow the obtention of peptide profiles resulting from an extensive protein hydrolysis. Furthermore, free amino acids are also released from the N- and C-terminals through the action of exopeptidases and, consequently, the remaining peptides are progressively reduced in size. Peptidases are commonly found in microorganisms. For instance, lactic acid bacteria contains an extracellular serin proteinase and several intracellular peptidases. In fact, many intracellular exopeptidases have been reported in the literature like the general aminopeptidase PepN in *L. Helveticus* and *L. sakei*, the glutamyl (aspartyl) specific aminopeptidase, PepA in

Streptococcus cremoris, *Lactococcus lactis* sp. and *Lb. delbrueckii* ssp. *lactis*, the proline specific peptidases, such as PepX and PepP in *Lactococcus lactis* ssp. *lactis*, X-prolyl di-peptidyl peptidase activity in *Leuconostoc mesenteroide*, *L. curvatus* and in *L. sakei*, di-peptidyl peptidases in *L. paracasei*, dipeptidase in *L. sakei*, *L. helveticus*, *L. plantarum*, *L. brevis*, *L. paracasei* and *L. casei* sp *casei*, arginyl aminopeptidase and tripeptidase in *L. sakei* (Bintsis et al., 2004; Macedo et al., 2010; Zotta et al., 2007; Stressler, González et al., 2010; Eisele, Schlayer, Lutz-Wahl & Fischer, 2013a; Stressler Eisele, Schlayer & Fischer, 2012; Stressler Eisele, Kranz & Fischer, 2014; Stressler et al., 2016; Flores & Toldrá, 2011). The yeast *Debaryomyces hansenii*.also contains endopeptidases like protease A and D and intracellular exopeptidases like prolyl and arginyl aminopeptidases (Santos, Santos-Mendonça, Sanz, Bolumar, Aristoy & Toldrá, 2001), all of them reported in Endopeptidases like neutral and alkaline protease and exopeptidases like X-prolyl di-peptidyl peptidase, leucine aminopeptidase, and dipeptidyl peptidases (DPP) IV and V have been reported in molds like *Aspergillus oryzae* and DPP V in *Aspergillus fumigatus* (Matsushita-Morita et al., 2011; Stressler et al., 2016).

4. Endogenous protein hydrolysis in foods

Endogenous food peptidases may be responsible for the release of polypeptides and bioactive peptides. The first step of proteolysis is the breakdown of proteins by endopeptidases into major fragments. Figure 3 shows how ubiquitin 60S ribosomal protein, a muscle protein, is degraded by endogenous muscle endo-peptidases into major fragments at cleaving sites Leu-Glu, Lys-Glu, Leu-Ile and Leu-Ser during the processing of dry-cured ham (Mora, Gallego, Aristoy, Fraser & Toldrá, 2015). In the case of short term processed foods, like fermented sausages, additional peptidases from different

254 sources such as certain lactic acid bacteria, yeasts or molds are needed for the
 255 generation of bioactive peptides. The extent of proteolysis can be confirmed after
 256 comparing the chromatographic profiles of the controls with those of the inoculated
 257 microorganism.

258 Dipeptides may be generated in foods through the action of di-peptidyl peptidases
 259 (DPP). So, such activity in *L. paracasei* is able to release dipeptides like Ala-Phe, Pro-
 260 Leu, Lys-Leu, Leu-Gly and Lys-Phe (Bintsis, Vafopoulou-Mastrojiannaki, Litopoulou-
 261 Tzanetaki, 2004), X-prolyl di-peptidyl peptidase activity releases particular proline-
 262 containing dipeptides in *Leuconostoc mesenteroides* and *L. curvatus* strains (Zotta,
 263 Ricciardi & Parente.2007), and DPP activity in *Leuconostoc mesenteroides*, releases
 264 Arg-Pro and Gly-Phe and additionally Gly-Pro in *L. paracasei* subsp *casei* (Macedo,
 265 Vieira, Poças & Malcata. 2010). Several dipeptides X.Pro and tripeptides X-Pro-Pro
 266 have been identified in casein hydrolysates with *Lb. helveticus* (Stressler, Eisele &
 267 Fischer, 2013).

268 Muscle foods contain endogenous muscle di-peptidyl peptidases, especially DPP I and
 269 DPP II, which are active at slightly acid pH (5.5-6.5) and are able to hydrolyse
 270 dipeptides like Ala-Gln, Arg-Gly, Asn-Pro, Ile-Leu, Ala-Gly, Ser-Gly, Ser-Gln located
 271 in the N-terminal. An example for the action of such di-peptidyl-peptidases is shown in
 272 figure 4 where dipeptide Pro-Ala is sequentially released from the N-terminal of myosin
 273 light chain I (Mora, Sentandreu & Toldrá, 2011). Proline and alanine are also released
 274 by the action of aminopeptidase activity. Muscle tri-peptidyl peptidase I is also active at
 275 slightly acid pH (5.5-6.5) and is able to release certain tripeptides like Ile-Ile-Pro, Arg-
 276 Gly-Ala, Gly-Asn-Pro, Gly-Ala-Gly, Gly-Pro-Gly located at the N-terminal (Mora,
 277 Gallego, Escudero, Reig, Aristoy & Toldrá, 2015).

278 Some of the released di-peptides might be further hydrolysed by di-peptidase activity
279 into their individual amino acids. This is especially relevant when those bioactive di-
280 peptides because the bioactivity would be lost when broken down and no beneficial
281 health effects would be observed. So, di-peptidase activity has been reported several
282 microorganisms like *L. plantarum* and *L. paracasei* that can hydrolyse Leu-Leu, Phe-
283 Ala, and also Ala-Phe, Tyr-Leu and Lys-Leu, at lower rate while other dipeptides like
284 Ala-Ala or Leu-Gly remain unaffected (González, Sacristán, Arenas, Fresno &
285 Tornadijo, 2010). *L. brevis* has higher di-peptidase activity on Leu-Leu, Tyr-Leu, Ala-
286 Ala, Leu-Gly, Ala-Phe, Lys-Leu and Phe-Ala and also *L. casei* sp *casei* but at much
287 lower rate (González et al 2010). Di-peptides are reported to be more efficiently taken
288 up by cellular transport systems and peptidases in *L. sakei* (Sinz & Schwab, 2012).
289 The released tri-peptides may be also hydrolysed into a single amino acid and a di-
290 peptide. As mentioned for di-peptides, this would be also damaging if those tri-peptides
291 are bioactive. A tri-peptidase from *L. sakei* was reported although tripeptides are also
292 readily cleaved by Pep N of a variety of lactic acid bacteria (Flores & Toldrá, 2011).
293 High aminopeptidase activity has been reported for *Leuconostoc mesenteroides* and *L.*
294 *curvatus* while *L. plantarum*, *L. pentosus* and *Weissella cibaria* showed a variable
295 enzymatic activity between strains (Zotta et al., 2007). In general, lactic acid bacteria
296 show aminopeptidase activity being able to release different amino acids from the N-
297 terminal. So, *L. plantarum*, *L. brevis* and *L. casei subsp casei* have been reported to
298 release alanine, lysine, proline and leucine (Herrerros et al., 2003), *L. paracasei subesp*
299 *casei* releases alanine, arginine, lysine, methionine and leucine (Bintsis et al., 2003;
300 Macedo et al., 2010), *L. sakei* releases alanine and leucine, *L. plantarum* releases
301 leucine and *L. paracasei subsp paracasei* releases alanine, lysine, proline and leucine
302 (González et al., 2010; Macedo et al., 2010). The yeast *Debaryomyces hansenii* was

reported to hydrolyse sarcoplasmic proteins and generate large amounts of most amino acids (Santos et al., 2001).

On the other hand, very low or negligible carboxypeptidase activity has been reported in cell-free extracts of several lactic acid bacteria (González et al., 2010; Herreros, et al., 2003), and a low activity for *L. paracasei subsp paracasei* to release phenylalanine and arginine (Bintsis et al., 2003; Macedo et al., 2010). However, endogenous carboxypeptidase activity is more evident in muscle-based foods where the presence of hydrophobic amino acids like phenylalanine, tyrosine, tryptophan, methionine, isoleucine, leucine, valine and proline residues in the C-terminal promotes its hydrolysis by endogenous muscle carboxypeptidase A. The rest of amino acids are preferentially hydrolysed by muscle carboxypeptidase B (Mora et al., 2015a).

The extent of proteolysis and the amount of generated bioactive peptides depends on multiple variables including the raw materials, the type of enzyme activity, the microbial population, and processing conditions. A first insight on small peptides generated in a model fermented sausage inoculated with *Lactobacillus curvatus* CRL705 and *Staphylococcus vitulinus* GV318 gave some information on potential routes for proteolysis during fermentation and ripening (López et al., 2015). Bioactive peptides were generated in dry-sausages containing added sodium caseinate and fermented with *Lactobacillus pentosus* and *Staphylococcus carnosus* (Mora et al., 2015b). In both cases, Staphylococci peptidases might be involved in peptide generation because CNS have been reported to exert an important proteolytic activity against meat proteins (Mauriello, Casaburi, Blaiotta & Villani, 2004). In addition, *L. pentosus* and *S. carnosus* have a proteinase attached to the cell wall that supports the extracellular casein degradation into oligopeptides that can be further eluted into the cytoplasm and be degraded by intracellular peptidases into smaller peptides and free amino acids (Chaves-

López et al., 2014). β -casein has been reported to be more hydrolysed than other types of caseins probably due to its abundance in proline, leucine and valine residues which are preferred by aminopeptidases and carboxypeptidases (Mora et al., 2015b).

Two hexapeptides with relevant antioxidant activity were isolated and identified after the simulated gastrointestinal digestion of Stracchino which is a soft cheese produced in the Northern Italy (Pepe et al., 2016).

The peptide profiles of nine months dry-cured ham after fractionation by gel filtration are shown in Figure 5 for ACE inhibitory activity and antioxidant activity measured through the DPPH and ferric reducing power. It can be observed that all 3 activities are concentrated in similar fractions corresponding to small peptides, with size <2500 Da.

5. Hydrolysis of food proteins by peptidases

Alternatively, proteins may be isolated and hydrolysed by specific commercial peptidases which can be obtained from different origins as listed in table 1. The hydrolysis is carried out in a reactor followed by separation/purification operations. The cleavage site of food proteins is very relevant and changes for each enzyme. For instance, trypsin may cleave proteins at the carboxy side of arginine and lysine residues, chymotrypsin cleaves on the carboxy side of aromatic or hydrophobic amino acids. Pepsin A prefers phenylalanine, leucine or glutamic acid at the C-terminal. Alcalase prefers the carboxy side of hydrophobic residues.

The progress of protein hydrolysis is usually followed with the degree of hydrolysis (DH). Figure 5 is showing the progress of hydrolysis of thornback ray muscle hydrolysate treated with Alcalase, Neutrase, an enzyme preparation from *Bacillus subtilis* A26 and an extract of crude alkaline proteases from *Raja clavata* (Lassoued et al., 2015a). So, the hydrolysis of a food protein with different peptidases will result in

different peptides patterns. Furthermore, the number or amount of released peptides as such is not the final target which must be focused on the number and amount of bioactive peptides. This was clearly reported for muscle hydrolysates treated with an extract of crude alkaline proteases and Neutrase, although not showing the highest DH (see figure 5) were reported as the most powerful to prevent DNA oxidation (Lassoued et al., 2015a). However, a similar study with Thornback ray gelatin showed that the hydrolysate treated with Alcalase was the most protective against DNA oxidation (Lassoued et al., 2015b). Similarly, lentil protein concentrates that were hydrolysed with Alcalase gave the highest yield of peptides even though the hydrolysis with Savinase gave more bioactive peptides (García-Mora, Peñas, Frías & Martínez-Villaluenga, 2014). So, the choice of the most adequate peptidase for each type of protein and target peptide bioactivity must be carefully studied and considered.

Commercial peptidases are sometimes not clearly defined in the manufacturers specifications and this may affect the degree of hydrolysis and its content in small peptides and free amino acids. In addition to the main enzyme activity, some side activities may be found (see Table 1). A good example is Flavourzyme, a peptidase extracted from *Aspergillus oryzae*, that was recently subjected to a nine step purification and characterization. The results showed the activity of 3 endopeptidases but also other enzymes like 2 aminopeptidases, 2 dipeptidylpeptidases and one amylase (Merz et al., 2015). Further, a characterisation of 10 commercial peptidases was performed through a three-step methodology (Merz, Claaßen, Appel, Berends, Rabe, Blank et al., 2016).

Exopeptidase activity, based on the release of free amino acids, was found in Alcalase 2.4L (Novozymes), Maxazyme NNP DS (DSM), Flavourzyme 1000L (Novozymes) and Protease AN (Amano Enzyme Inc.). Such exopeptidase activity can be attributed to aminopeptidase and carboxypeptidase activity. The rest of assayed commercial

peptidases were Biopraxe SP-20FG (Nagase), Collupulin 200 L (DSM), Corolase2TS (AB Enzymes), Promod 439 L (Biocatalysts Ltd.), Proteinase T (DuPont) and Protin SD-AY10 (Amano Enzyme Inc.) and they were reported to exert majorly endopeptidase activity so that low degree of hydrolysis and poor generation of free amino acids may be expected (Merz et al., 2016). Other enzymes have shown also different peptide patterns. For instance, Neutrase was reported to give shorter peptide fragments than papain when hydrolysing rawhide collagen (Damrongsakkul, Ratanathammapan, Komolpis & Tanthapanichakoon, 2008). Whey protein concentrates hydrolysed with Neutrase also gave better iron absorption than those with papain or Alcalase (Ou et al., 2010). Pepsin, trypsin, protease M and flavourzyme have been successfully tested to produce calcium chelating peptides from different food protein sources (Sun, Wu, Du, Tang, Liu & Fu, 2016).

Sequential hydrolysis with different peptidase preparations may be used to produce bioactive peptides of interest. For instance, the hydrolysis of hen egg white lysozyme combining trypsin and papain gave a better yield of antioxidant and antimicrobial peptides than the use of trypsin or papain alone (Memarpoor-Yazdi, Asoodeh & Chamani, 2012). *Brassica carinata* proteins were sequentially hydrolysed with immobilised trypsin, chymotrypsin, and carboxypeptidase A and resulted in an enriched fraction with antioxidant peptides (Pedroche et al., 2007). Smooth hound viscera from *M. mustelus* was hydrolysed using commercial proteases (Purafect, Neutrase and Esperase) and combinations of such commercial enzymes with endogenous enzymes, being the last one the best option for the higher recovery of antioxidant, ACE-inhibitory and antibacterial peptides (Abdelhedi et al., 2016). Eight commercial enzyme preparations were combined and used to obtain bioactive peptides from protein hydrolysates of defatted salmon backbone. The highest antioxidant and ACE inhibitory

activity was obtained with trypsin, bromelain, papain and protamex treatment (Slizyte, Rommi, Mozuraityte, Eck, Five & Rustad, 2016). Other authors reported an original way to improve the peptide profile and its bioactivity in a protein hydrolysate (Xu, Kong & Zhao, 2014). This was achieved with plastein that has the ability to reverse the hydrolytic action by peptidases, forming polypeptides. So, casein was first hydrolysed with Neutrase to generate ACE inhibitory peptides and this hydrolysate was then used as substrate for further plastein reaction that once optimised could increase the ACE inhibitory activity of the hydrolysate (Xu et al., 2014).

Some caution must be taken when using commercial enzymes especially in the efficacy and reproducibility of protein hydrolysis and also the enzymes stability. Some batch to batch variability may be observed due to variations in the activity of certain enzymes. For instance, flavorzyme was reported to have some variability in casein hydrolysis that was attributed to loss of endopeptidase activity along the storage time (Merz, Appel, Berends, Rabe & Blank, 2016). In the case of endogenous hydrolysis, the peptide profile may also change for similar types of foods due to different raw materials that may have different endogenous enzymes profiles but also to changes in processing.

6. Conclusions

The final result of protein hydrolysates consists of a pool of peptides with different sequences and lengths, some of them with a relevant bioactivity depending on the particular food and type and conditions of hydrolysis. Thus, the generated small peptides may exhibit a wide range of bioactivities such as angiotensin converting enzyme (ACE) inhibitory activity, antioxidant, antithrombotic, hypoglucemic, hypocholesterolemic, and antimicrobial activity among others. However, it must be considered that the generated bioactive peptides, either endogenously in food or a

protein hydrolysate, may be further hydrolysed when ingested through the gastrointestinal digestion. Further, those peptides should be absorbed intact through the intestinal barrier and reach the blood stream to exert their physiological action, overcoming the potential sequence modifications by brush border peptidases during transepithelial transport.

Acknowledgements

The research leading to these results received funding from Grant AGL2014-57367-R from the Spanish Ministry of Economy, Industry and Competitiveness and FEDER funds and from the European Union 7th Framework Programme (FP7/2007-2013) under Grant Agreement 312090 (BACCHUS). This publication reflects only the author views and the Community is not liable for any use made of the information contained therein. The Juan de la Cierva postdoctoral contract to LM is also acknowledged. Mass spectrometry analysis was performed in the SCSIE_University of Valencia Proteomics Unit, a member of ISCIII ProteoRed Proteomics Platform.

Conflicts of interest

All authors of this manuscript declare that they do not have any conflict of interest.

References

1. Abdelhedi, O., Jridi, M., Jemil, I., Mora, L., Toldrá, F., Aristoy, M.C., Boualga, A., Nasri, M., Nasri, M. (2016) Combined biocatalytic conversion of smooth hound viscera: Protein hydrolysates elaboration and assessment of their

antioxidant, anti-ACE and antibacterial activities. *Food Research International*,
86, 9–23.

2. Bintsis, T., Vafopoulou-Mastrogiannaki, A., Litopoulou-Tzanetaki, E.,
Robinson, R.K. (2003) Protease, peptidase and esterase activities by lactobacilli
and yeast isolates from Feta cheese brine. *Journal of Applied Microbiology*, 95,
68–77
3. Capriotti, A. L., Caruso, G., Cavaliere, C., Samperi, R., Ventura, S., Chiozzi, R.
Z., Laganà, A. (2015). Identification of potential bioactive peptides generated by
simulated gastrointestinal digestion of soybean seeds and soy milk proteins.
Journal of Food Composition and Analysis, 44, 205-213.
4. Castellano, P., Mora, L., Escudero, E., Vignolo, G., Aznar, R., & Toldrá, F.
(2016). Antilisterial peptides from Spanish dry-cured hams: purification and
identification. *Food Microbiology*, 59, 133-141.
5. Charoenphun, N., Cheirsilp, B., Sirinupong, N., Youravong W. (2013) Calcium-
binding peptides derived from tilapia (*Oreochromis niloticus*) protein
hydrolysate. *European Food Research and Technology*, 236, 57–63.
6. Chaves-López, C., Serio, A., Paparella, A., Martuscelli, M., Corsetti, A., Tofalo,
R., Suzzi, G. (2014) Impact of microbial cultures on proteolysis and release of
bioactive peptides in fermented milk. *Food Microbiology*, 42, 117–121.
7. Choi, D.W., Lee, J.H., Chun, H.H., Song K.B., (2012) Isolation of a calcium-
binding peptide from bovine serum protein hydrolysates. *Food Science and
Biotechnology*, 21, 1663–1667.
8. Corrêa, A. P. F., Daroit, D. J., Fontoura, R., Meira, S. M. M., Segalin, J.,
Brandelli, A. (2014). Hydrolysates of sheep cheese whey as a source of

478 bioactive peptides with antioxidant and angiotensin-converting enzyme
 479 inhibitory activities. *Peptides*, 6, 48-55.

480 9. De Oliveira CF, Corrêa AP, Coletto D, Daroit DJ, Cladera-Olivera F, Brandelli
 481 A. (2015). Soy protein hydrolysis with microbial protease to improve
 482 antioxidant and functional properties. *Journal of Food Science & Technology*,
 483 52, 2668-78.

484 10. Dong-Eun Sung, Jeongok Lee, Youngshin Han, Dong-Hwa Shon, Kangmo Ahn,
 485 Sangsuk Oh, Jeong-Ryong Do. (2014) Effects of enzymatic hydrolysis of
 486 buckwheat protein on antigenicity and allergenicity. *Nutrition Research and
 487 Practice*, 8, 278-283.

488 11. Escudero, E., Mora, L., & Toldrá, F. (2014). Stability of ACE inhibitory ham
 489 peptides against heat treatment and in vitro digestion. *Food Chemistry*, 161,
 490 305-311.

491 12. Escudero, E., Sentandreu, M.A., Arihara, K., Toldrá, F. (2010) Angiotensin I
 492 converting enzyme inhibitory peptides generated from in vitro gastrointestinal
 493 digestion of pork meat. *Journal of Agricultural & Food Chemistry*, 58, 2895-
 494 2901.

495 13. Fernández, M., Benito, M. H., Martín, A., Casquete, R., Córdoba, J. J., & Córdoba,
 496 M. G. (2016). Influence of starter culture and a protease on the generation of ACE-
 497 inhibitory and antioxidant bioactive nitrogen compounds in Iberian dry-fermented
 498 sausage ‘salchichón’. *Heliyon*, 31, 2 (3).

499 14. Ferraro, V., Carvalho, A.P., Piccirillo, C., Santos, M.M., Castro, P.M.L., Pintado,
 500 M.E. (2013). Extraction of high added value biological compounds from sardine,
 501 sardine-type fish and mackerel canning residues. *Materials Science and
 502 Engineering C*, 33, 3111-20.

15. Fitzgerald R.J., Meisel H. (2000) Milk protein-derived peptide inhibitors of angiotensin-I-converting enzyme. *British Journal of Nutrition*, 84, S33–S37.
16. Flores, M., Toldrá, F. (2011) Microbial enzymatic activity for improved fermented meats. *Trends in Food Science and Technology*, 22, 81-90
17. Freiding, S., Gutsche, K.A., Ehrmann, M.A., Vogel, R. (2011) Genetic screening of *Lactobacillus sakei* and *Lactobacillus curvatus* strains for their peptidolytic system and amino acid metabolism, and comparison of their volatiles in a model system. *Systematic and Applied Microbiology* 34, 311–320.
18. Gallego, M., Mora, L., Aristoy, M.C., Toldrá, F. (2015) Titin-derived peptides as processing time markers in dry-cured ham. *Food Chemistry*, 167, 326-339.
19. García-Mora, P., Peñas, E., Frias, J., Martínez-Villaluenga, C. (2014) Savinase, the Most Suitable Enzyme for Releasing Peptides from Lentil (*Lens culinaris* var. Castellana) Protein Concentrates with Multifunctional Properties. *Journal of Agricultural and Food Chemistry*, 62, 4166–4174.
20. Georgieva, D.N., Stoeva, S., Voelter, W., Genov, N., Betzel, C. (2001) Substrate specificity of the highly alkalophilic bacterial proteinase Esperase: Relation to the X-ray structure. *Current Microbiology*, 42, 368-371.
21. González, L., Sacristán, N., Arenas, R., Fresno, J.M., Tornadijo, E. (2010) Enzymatic activity of lactic acid bacteria (with antimicrobial properties) isolated from a traditional Spanish cheese. *Food Microbiology*, 27, 592-597.
22. Gu, Y., Majumder, K., & Wu, J. (2011). QSAR-aided in silico approach in evaluation of food proteins as precursors of ACE inhibitory peptides. *Food Research International*, 44, 2465-2474.

- 526 23. Guesdon, B., Pichon, L., Tomé, D. (2006) Opioid peptides. In: *Nutraceutical*
527 *Proteins and Peptides in Health and Disease*, (Y. Mine and F. Shahidi, Eds.),
528 Francis and Taylor Group, Boca Raton, Fl, pp. 367-376.
- 529 24. Guo, L., Harnedy, P.A., O'Keeffe, M.B., Zhang, L., Li, B., Hou, H., Fitzgerald,
530 R.J. (2015) Fractionation and identification of Alaska pollock skin collagen-
531 derived mineral chelating peptides. *Food Chemistry*, 173, 536–542.
- 532 25. Herreros, M.A., Fresno, J.M., GonzálezPrieto, M.J.,
533 Tornadijo, M.E., 2003. Technological characterization of lactic acid bacteria
534 isolated from Armada cheese (a Spanish goats' milk cheese). *International Dairy*
535 *Journal* 13, 469-479.
- 536 26. Huang, D., Ou, B., Prior, R.L. (2005). The chemistry behind antioxidant
537 capacity assays. *Journal of Agricultural and Food Chemistry*, 53, 1841-1856.
- 538 27. Huang, S.L., Zhao, L.N., Cai, X., Wang, S.Y., Huang, Y.F., Hong, J., Rao, P.F.
539 (2015) Purification and characterisation of a glutamic acid-containing peptide
540 with calcium-binding capacity from whey protein hydrolysate. *Journal of Dairy*
541 *Research*, 82, 29–35.
- 542 28. Ji, N., Sun, C. X., Zhao, Y. X., Xiong, L., Sun, Q. J. (2014). Purification and
543 identification of antioxidant peptides from peanut protein isolate hydrolysates
544 using UHR-Q-TOF mass spectrometer. *Food Chemistry*, 161, 148–154.
- 545 29. Kitchener, R.L., Grunden, A.M. (2012) Prolidase function in proline metabolism
546 and its medical and biotechnological applications. *Journal of Applied*
547 *Microbiology*, 113, 233–247.
- 548 30. Lassoued, I., Mora, L., Nasri, R., Aydi, M., Toldrá, F., Aristoy, M.C., Barkia,
549 A., Nasri, M. (2015a) Characterization, antioxidative and ACE inhibitory

- 550 properties of hydrolysates obtained from Thornback Ray (*Raja clavata*) muscle.
551 *Journal of Proteomics*, 128, 458-468.
- 552 31. Lassoued, I., Mora, L., Nasri, R., Jridi, M., Toldrá, F., Aristoy, M.C., Barkia, A.,
553 Nasri, M. (2015b) Characterization and comparative assessment of antioxidant
554 and ACE inhibitory activities of thornback ray gelatin hydrolysates. *Journal of*
555 *Functional Foods* 13, 225–238.
- 556 32. Li G.H., Le G.W., Shi Y.H., Shrestha S. (2004) Angiotensin I–Converting
557 enzyme inhibitory peptides derived from food proteins and their physiological
558 and pharmacological effects. *Nutrition Research*, 24, 469–486
- 559 33. Li-Chan, ECY (2015). Bioactive peptides and protein hydrolysates:research
560 trends and challenges for application as nutraceuticals and functional food
561 ingredients. *Current Opinion in Food Science*, 1, 28-37.
- 562 34. Liu, R., Xing, L., Fu, Q., Zhou, G., and Zhang, W. (2016) A Review of
563 Antioxidant Peptides Derived from Meat Muscle and By-Products. *Antioxidants*
564 (Basel), 5, 32.
- 565 35. López, C.M., Bru, E., Vignolo, G., Fadda, S.G. (2015) Identification of small
566 peptides arising from hydrolysis of meat proteins in dry fermented sausages.
567 *Meat Science*, 104, 20-29.
- 568 36. Lv, Y., Bao, X., Liu, H., Ren, J., Guo S. (2013) Purification and characterization
569 of calcium-binding soybean protein hydrolysates by Ca²⁺/Fe³⁺ immobilized
570 metal affinity chromatography (IMAC). *Food Chemistry*, 141, 1645–1650.
- 571 37. Macedo, A.C., Vieira, M., Poças, R., Malcata, F.X. (2010) Peptide hydrolase
572 system of lactic acid bacteria isolated from Serra da Estrela cheese. *International*
573 *Dairy Journal*, 10, 769-774.

38. Matsushita-Morita, M., Tada, S., Suzuki, S., Hattori, R., Marui, J., Furukawa, I., Yamagata, Y., Amano, H., Ishida, H., Takeuchi, M., Kashiwagi, Y., Kusumoto, K.I. (2011) Overexpression and Characterization of an Extracellular Leucine Aminopeptidase from *Aspergillus oryzae*. *Current Microbiology*, 62, 557 -564.
39. Mauriello, G., Casaburi, A., Blaiotta, G., Villani, F. (2004) Isolation and technological properties of coagulase negative staphylococci from fermented sausages of Southern Italy. *Meat Science*, 67, 149-158.
40. MacDonald-Wicks, L.K., Wood, L.G., Garg, M.L. (2006). Methodology for the determination of biological antioxidant capacity in vitro: a review. *Journal of the Science of Food and Agriculture*, 86, 2046-2056.
41. Meinlschmidt, P., Sussmann, D., Schweiggert-Weisz, U., Eisner, P. (2016) Enzymatic treatment of soy protein isolates: effects on the potential allergenicity, technofunctionality, and sensory properties. *Food Science & Nutrition*, 4, 11–23.
42. Memarpoor-Yazdi, M., Asoodeh, A., Chamani, J. (2012) A novel antioxidant and antimicrobial peptide from hen egg white lysozyme hydrolysates. *Journal of Functional Foods*, 4, 278-286.
43. Merz, M., Appel, D., Berends, P., Rabe, S., Blank, I., Stressler, T., Fischer, L. (2016) Batch-to-batch variation and storage stability of the commercial peptidase preparation Flavourzyme in respect of key enzyme activities and its influence on process reproducibility. *European Food Research and Technology*, 242, 1005–1012.
44. Merz, M., Eisele, T., Berends, P., Appel, D., Rabe, S., Blank, I., Stressler, T., Fischer, L. (2015) Flavourzyme, an enzyme preparation with industrial

598 relevance: automated nine-step purification and partial characterization of eight
 599 enzymes. *Journal of Agricultural and Food Chemistry*, 63, 5682 -5693.

600 45. Merz, M., Claaßena, W., Appel, D., Berends, P., Rabe, S., Blank, I., Stressler,
 601 T., Fischer, L. (2016) Characterization of commercially available peptidases in
 602 respect of the production of protein hydrolysates with defined compositions
 603 using a three-step methodology. *Journal of Molecular Catalysis. B: Enzymatic*,
 604 127, 1-10.

605 46. Mihulová M., Vejlupeková M., Hanušová J., Štětina J., Panovská Z. (2013)
 606 Effect of modified whey proteins on texture and sensory quality of processed
 607 cheese. *Czech Journal of Food Science*, 31, 55--558.

608 47. Mohanty, D.P., Mohapatra, S., Misra, S., Sahu, P. S. (2016). Milk derived
 609 bioactive peptides and their impact on human health – A review. *Saudi Journal*
 610 *of Biological Sciences*, 23, 577-583.

611 48. Mora, L., Escudero, E., Arihara, K. & Toldrá, F. (2015) Antihypertensive effect
 612 of peptides naturally generated during Iberian dry-cured ham processing. *Food*
 613 *Research International*, 78, 71-78.

614 49. Mora, L., Gallego, M., Escudero, E., Reig, M., Aristoy, M-C., Toldrá, F. (2015a)
 615 Small peptides hydrolysis in dry-cured meats. *International Journal of Food*
 616 *Microbiology*, 212, 9-15.

617 50. Mora, L., Gallego, M., Aristoy, M.C., Fraser, P., Toldrá, F. (2015) Peptides
 618 naturally generated from ubiquitin-60S ribosomal protein as potential
 619 biomarkers of dry-cured ham processing time. *Food Control*, 48, 102-107.

620 51. Mora, L., Gallego, M., Aristoy, M-C., Toldrá, F. (2015b) A peptidomic
 621 approach to study the contribution of added casein proteins to the peptides

- profile in Spanish dry-fermented sausages. *International Journal of Food Microbiology*, 212, 41-48.
52. Mora, L., Sentandreu, M.A., Toldrá, F. (2011) Intense degradation of myosin light chain isoforms after dry-cured ham processing. *Journal of Agricultural & Food Chemistry*, 2011, 59, 3884-3892.
53. Mora, L., Reig, M., Toldrá, F. (2014) Bioactive peptides generated from meat industry by-products. *Food Research International*, 65, 344-349.
54. Nongonierma, A.B. and FitzGerald, R.J. (2016) Strategies for the discovery, identification and validation of milk protein-derived bioactive peptides. *Trends in Food Science & Technology*, 50, 26-43
55. Oseguera-Toledo, ME, González de Mejía, E., Reynoso-Camacho, R., Cardador-Martínez, A., Amaya-Llano, S.L. (2014). Proteins and bioactive peptides: Mechanisms of action on diabetes management. *Nutrafoods*, 13, 147-157.
56. Ou, K., Liu, Y., Zhang, L., Yang, X., Huang, Z., Nout, M.J.R., Liang, J. (2010) Effect of Neutrase, Alcalase, and Papain Hydrolysis of Whey Protein Concentrates on Iron Uptake by Caco-2 Cells. *Journal of Agricultural and Food Chemistry*, 58, 4894-4900.
57. Pedroche, J., Yust, M. M., Lqari, H., Megias, C., Girón-Calle, J. Alaiz, M., Vioque J., Millán, F. (2007). Obtaining of *Brassica carinata* protein hydrolysates enriched in bioactive peptides using immobilized digestive proteases. *Food Research International*, 40, 931-938.
58. Pepe, G., Sommella, E., Ventre, G., Scala, M.C., Adesso, S., Ostacolo, C., Marzocco, S., Novellino, E., Campiglia, P. (2016) Antioxidant peptides released from gastrointestinal digestion of “Stracchino” soft cheese: Characterization, in

647 vitro intestinal protection and bioavailability. *Journal of Functional Foods*, 26,
648 494 -505.

649 59. Pihlanto-Leppälä, A. (2001). Bioactive peptides derived from bovine whey
650 proteins: Opioid and ACE-inhibitory peptides. *Trends in Food Science and*
651 *Technology*, 1, 347-356.

652 60. Rajapakse N., Mendis E., Jung W.K., Je J.Y., Kim S.K. (2005). Purification of a
653 radical scavenging peptide from fermented mussel sauce and its antioxidant
654 properties. *Food Research International*, 38, 175–182.

655 61. Ranathunga S., Rajapakse N., Kim S.-K. (2006) Purification and
656 characterization of antioxidative peptide derived from muscle of conger eel
657 (Conger myriaster). *European Food Research and Technology*, 222, 310–315.

658 62. Reyes-Díaz, González-Córdova, Hernández-Mendoza, Vallejo-Córdoba, (2016)
659 Immuno-modulating peptides obtained from milk proteins. *Interciencia*, 41,
660 84-91.

661 63. Ryder, K., Bekhit, A. E. D., McConnell, M., Carne, A. (2016). Towards
662 generation of bioactive peptides from meat industry waste proteins: Generation
663 of peptides using commercial microbial proteases, *Food Chemistry*, 208, 42-50.

664 64. Santos, N.N., Santos-Mendonça, C., Sanz, Y., Bolumar, T., Aristoy, M.C.,
665 Toldrá, F. (2001) Hydrolysis of pork muscle sarcoplasmic proteins by
666 *Debaryomyces hansenii*. *International Journal of Food Microbiology*, 68, 199-
667 206

668 65. Sentandreu, M.A., Toldrá, F. (2000) Purification and biochemical properties of
669 dipeptidyl peptidase I from porcine skeletal muscle. *Journal of Agricultural and*
670 *Food Chemistry*, 48, 5014-5022.

66. Sentandreu, M.A., Toldrá, F. (2001) Partial purification and characterisation of dipeptidyl peptidase II from porcine skeletal muscle. *Meat Science*, 57, 93-103.
67. Sinz, Q., Schwab, W. (2012) Metabolism of amino acids, dipeptides and tetrapeptides by *Lactobacillus sakei*. *Food Microbiology*, 29, 215-223.
68. Slizyte, R., Rommi, K., Mozuraityte, R., Eck, P., Five, K., Rustad, T. (2016) Bioactivities of fish protein hydrolysates from defatted salmon backbones. *Biotechnology Reports*, 11, 99–109.
69. Stefanucci, A., Mollica, A., Macedonio, G., Zengin, G., Ahmed, A.A. & Novellino, E. (2017) Exogenous Opioid Peptides Derived From Food Proteins and Their Possible Uses as Dietary Supplements: A Critical Review. *Food Reviews International*, DOI: 10.1080/87559129.2016.1225220
70. Stressler, T., Eisele, T., Kranz, B., Fischer, L. (2014) PepX from *Lactobacillus helveticus*: Automated multi-step purification and determination of kinetic parameters with original tripeptide substrates. *Journal of Molecular Catalysis. B: Enzymatic*, 108, 103 -110.
71. Stressler, T., Eisele, T., Schlayer, M., Fischer, L. (2012) Production, active staining and gas chromatography assay analysis of recombinant aminopeptidase P from *Lactococcus lactis* ssp. *lactis* DSM 20481. *AMB Express*, 2, 39.
72. Stressler, T., Eisele, T., Schlayer, M., Lutz-Wahl, S., Fischer, L. (2013a) Characterization of the Recombinant Exopeptidases PepX and PepN from *Lactobacillus helveticus* ATCC 12046 Important for Food Protein Hydrolysis. *PLoS ONE*, 8, 70055.
73. Stressler, T., Eisele, T., Fischer, L. (2013b) Simultaneous monitoring of twelve angiotensin I converting enzyme inhibitory peptides during enzymatic β -casein

- 695 hydrolysis using *Lactobacillus* peptidases. *International Dairy Journal*, 30, 96-
696 102.
- 697 74. Stressler, T., Eisele, T., Fischer, L. (2013) Simultaneous monitoring of twelve
698 angiotensin I converting enzyme inhibitory peptides during enzymatic beta-
699 casein hydrolysis using *Lactobacillus* peptidases. *International Dairy Journal*,
700 30, 96–102.
- 701 75. Suetsuna, K., Ukeda, H., Ochi, H. (2000). Isolation and characterization of free
702 radical scavenging activities peptides derived from casein. *Journal of*
703 *Nutritional Biochemistry*, 11, 128–131.
- 704 76. Tanzadehpanah, H., Asoodeh, A., Chamani, J. (2012). An antioxidant peptide
705 derived from Ostrich (*Struthio camelus*) egg white protein hydrolysates. *Food*
706 *Research International*, 49, 105–111.
- 707 77. Vercruysse, L., Van Camp, J., Smagghe, G. (2005) ACE inhibitory peptides
708 derived from enzymatic hydrolysates of animal muscle protein: A review.
709 *Journal of Agricultural and Food Chemistry*, 53, 8106-8115.
- 710 78. Xu, W., Kong, B.H., Zhao, X.H. (2014) Optimization of some conditions of
711 Neutrase-catalyzed plastein reaction to mediate ACE-inhibitory activity in vitro
712 of casein hydrolysate prepared by Neutrase. *Journal of Food Science and*
713 *Technology-Mysore*, 51, 276-284.
- 714 79. Yoshida, N., Tsuruyama, S., Nagata, K., Hirayama, K., Noda, K., Makisumi, S.
715 (1988) Purification and characterization of an acidic amino acid specific
716 endopeptidase of *Streptomyces griseus* obtained from a commercial preparation
717 (Pronase). *Journal of Biochemistry*, 104, 451-6.

80. Zotta, T., Ricciardi, A. & Parente, E. (2007) Enzymatic activities of lactic acid bacteria isolated from Cornetto di Matera sourdoughs. *International Journal of Food Microbiology*, 115, 165-172.

LEGENDS FOR THE FIGURES

Figure 1.- Scheme of protein hydrolysis and enzymes involved.

Figure 2.- Scheme of the generation of bioactive peptides from protein hydrolysis in foods and/or the hydrolysis of isolated food proteins.

Figure 3.- Peptides identified by nanoESI-LC-MS/MS derived from ubiquitin 60S ribosomal protein (UniProtKB/TrEMBL protein database accession number P63053). Endopeptidase activity is showed in black arrows. Adapted from Mora, Gallego, Aristoy, Fraser, Toldrá. 2015. Peptides naturally generated from ubiquitin-60S ribosomal protein as potential biomarkers of dry-cured ham processing time. *Food Control*, 48, 102-107.

Figure 4.- Intense degradation of Myosin Light Chain 1 (accession number A1XQT6_PIG in UniProtKB/TrEMBL database), evidencing the action of amino peptidases (in dark black) and dipeptidyl peptidases (in light black). Adapted from Mora, Sentandreu and Toldrá. 2011. Intense degradation of myosin light chain isoforms after dry-cured ham processing. *Journal of Agricultural & Food Chemistry*, 59, 3884-3892.

Figure 5.- Hydrolysis curves of thornback ray muscle hydrolysates (TRMHs) treated with Alcalase (TRMH-Alcalase), Neutrase (TRMH-Neutrase), enzyme preparation from *Bacillus subtilis* A26 (TRMH-A26) and crude alkaline protease extract from *R. clavata* (TRMH-Crude). Reproduced from Lassoued, Mora, Nssri, Aydi, Toldrá, Aristoy, Barkia and Nasri. 2015. Characterization, antioxidative and ACE inhibitory properties of hydrolysates obtained from Thornback Ray (*Raja clavata*) muscle. Journal of Proteomics, 128, 458-468, with permission from Elsevier.

Table 1.- Commercial enzyme preparations with specific characteristics and some relevant applications.

Commercial preparation	Origin	Manufacturers	Activity	Cleavage sites	Application	Literature
Flavourcyme 1000M	<i>Aspergillus oryzae</i>		3 endopeptidases 2 aminopeptidases 2 dipeptidylpeptidases 1 α -amylase		Cereals Calcium chelating peptides, Soy	Merz et al., 2015 Huang et al., 2015 Meinlschmidt et al, 2016
Valkerase	<i>Bacillus licheniformis</i>	Bri Enzymes	Keratinase, Serin Endopeptidase	Non especific	Feather meal	-
Prolidase	<i>L-lactis cremoris</i> Other many sources		Dipeptidase	Bonds including proline or hydroxyproline	Cheese making	Kitchener and Grunden, 2012
Biopraxe SP-20FG	<i>Bacillus sp</i>		Subtilisin Endo metalloprotease Aminopeptidase			Merz et al., 2015
Neutrase	<i>Bacillus subtilis</i> <i>B. amyloliquefaciens</i>	Novozymes	Metalloprotease		Collagen Calcium- and iron-chelating peptides Soy	Ou et al., 2010 Meinlschmidt et al, 2016
Alcalase 2.4 L	<i>Bacillus licheniformis</i>	Novozymes	Subtilisin Alkaline serin endopeptidase Extracellular neutral metallo protease Aminopeptidase	Non especific	Calcium-chelating peptides	Choi et al., 2012 Charoenphun et al., 2013

755 Figure 1

756

Proteolysis phenomena





Figure 1.- Scheme of protein hydrolysis and enzymes involved.

757

758



760 Figure 2.- Scheme of the generation of bioactive peptides from protein hydrolysis in foods and/or the hydrolysis of isolated food proteins

761

762

P ₀	Sequence		P ₁
3'	FVKTLTGKTITLEVEPSDTIENVKAKIQDKEGIPPDQQRLLIFAGKQLEDGRTLSDYNIQKESTL	67'	
I	FVKTLTGKTITL		E
L	EVEPSDTIENVKAKIQDK		E
K	EGIPPDQQRLL		I
L	IFAGKQLEDGRTL		S
L	SDYNIQKESTL		H

Figure 3. Peptides identified by nanoESI-LC-MS/MS derived from ubiquitin 60S ribosomal protein (UniProtKB/TrEMBL protein database accession number P63053). Endopeptidase activity is showed in black arrows. Adapted from Mora, L., Gallego, M., Aristoy, MC., Fraser, PD., Toldrà, F. (2015) Peptides naturally generated from ubiquitin-60S ribosomal protein as potential biomarkers of dry-cured ham processing time. Food Control, 48, 102-107.

764

765

766

Figure 4




Figure 4. Intense degradation of Myosin Light Chain 1 (accession number A1XQT6_PIG in UniProtKB/TrEMBL database), evidencing the action of amino peptidases (in dark black) and dipeptidyl peptidases (in light black).

Figure 5

